

Implications of structural diversity for seasonal and annual carbon dioxide fluxes in two temperate deciduous forests

Tamrakar, Rijan; Rayment, Mark B.; Moyano, Fernando; Mund, Martina; Knohl, Alexander

Agricultural and Forest Meteorology

DOI:

[10.1016/j.agrformet.2018.08.027](https://doi.org/10.1016/j.agrformet.2018.08.027)

Published: 15/12/2018

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Tamrakar, R., Rayment, M. B., Moyano, F., Mund, M., & Knohl, A. (2018). Implications of structural diversity for seasonal and annual carbon dioxide fluxes in two temperate deciduous forests. *Agricultural and Forest Meteorology*, 263, 465-476.
<https://doi.org/10.1016/j.agrformet.2018.08.027>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract

The effects of structural diversity on the carbon dioxide exchange (CO_2) of forests has become an important area of research for improving the predictability of future CO_2 budgets. We report the results of a paired eddy covariance tower experiment with 11 years of data on two forest sites of similar mean stand age, near-identical site conditions, and dominated by beech trees (*Fagus sylvatica*), but with a very different stand structure (incl. age, diameter distribution, stocks of dead wood and species composition) because of different management regimes. Here we address the question of how management and related structural diversity may affect CO_2 fluxes, and tested the hypothesis that more structurally diverse stands are less sensitive to variations in environmental and biotic drivers. Higher annual net ecosystem productivity (NEP) was observed in the homogenous, managed, and even-aged forest ($585 \pm 57.8 \text{ g C m}^{-2} \text{ yr}^{-1}$), than in the structurally diverse, unmanaged, and uneven-aged forest ($487 \pm 144 \text{ g C m}^{-2} \text{ yr}^{-1}$). About two-third of the difference in NEP between the sites was contributed by a higher annual gross primary productivity (GPP, 1627 ± 164 vs $1558 \pm 118 \text{ g C m}^{-2} \text{ yr}^{-1}$) and one-third by a lower annual ecosystem respiration (Reco, 1042 ± 60 vs $1071 \pm 96 \text{ g C m}^{-2} \text{ yr}^{-1}$) in the homogenous forest. Spring (April - May) and summer (June – July) were the two main seasons contributing to the overall annual differences between the sites, also, the sensitivities of seasonal NEP and GPP to environmental variables were stronger in the homogenous forest during those periods. Inter-annual variation of NEP was higher in the homogenous forest (Coefficient of variation (CV) = 25 %) compared to the heterogeneous forest (CV = 12 %). At annual time scale, the higher variability of NEP in the homogenous forest is mainly attributed to biotic factors such fruit production and a time-dependent growth trend, outweighing differences in environmental sensitivities.

Keywords: structural diversity; eddy covariance; temperate deciduous forest; CO_2 flux.

26 1. Introduction

27 The carbon uptake of forests is affected by changes in both environmental and biotic
28 factors (Chen et al., 2015; Ciais et al., 2005). The former includes temperature, radiation,
29 water and nutrient availability, and their intra and inter-annual variability. Biotic factors
30 include plant functional traits such as plant physiological parameters, nutrient status,
31 structure, phenology, etc., that govern photosynthesis and respiration process (Jensen et
32 al., 2017) as well as inter- and intra-specific competition. Identifying and understanding
33 the factors that contribute to the variability in net carbon dioxide (CO₂) uptake, i.e. net
34 ecosystem productivity (NEP), between forest ecosystems and the atmosphere is crucial
35 for understanding how forests will respond to and affect future climate (Baldocchi et al.,
36 2001; IGBP Terrestrial Carbon Working Group, 1998; Luo et al., 2015) as well as for
37 answering questions relevant to forest management and ecology.

38 Many eddy covariance (EC) flux studies (e.g. Barr et al., 2007; Dragoni et al., 2011; Hui
39 et al., 2003; Humphreys and Lafleur, 2011; Jensen et al., 2017; Kitamura et al., 2012;
40 Richardson et al., 2009; Shao et al., 2016, 2015; Wu et al., 2013; Yuan et al., 2009) have
41 attributed the inter-annual variability (IAV) of NEP variously to climatic variables, to
42 phenological changes induced by climatic variables and to biotic changes, with
43 Richardson et al. (2007) contending that, on an annual scale, variation in NEP is more
44 strongly dominated by changes in biotic factors than by climate. To date, most studies
45 have focused on understanding effects of climate and biotic changes on CO₂ fluxes at
46 single sites (Granier et al., 2008; Pilegaard et al., 2011; Wilkinson et al., 2012) or across
47 contrasting ecosystem types (Baldocchi and Xu, 2005; Chu et al., 2016; Jensen et al.,
48 2017; Ma et al., 2007; Novick et al., 2015; Pereira et al., 2007; Shao et al., 2016, 2015,
49 2014; Wu et al., 2012). Other studies have used multiple sites from across global and
50 regional networks to understand the variability of CO₂ fluxes from different plant
51 functional types and/or climatic zones (Beer et al., 2010; Chen et al., 2015; Law et al.,

2002; Musavi et al., 2017). Such studies have been beneficial for understanding the underlying causes of variability in CO₂ uptake, but because flux stations are not closely located, there are typically very large differences in the environmental conditions between sites, making it challenging to disentangle the effects of environmental vs biotic factors. The short period of time analysed is also a limitation found in some studies (Anthoni et al., 2004; Hommeltenberg et al., 2014; Jensen et al., 2017). Only a few have investigated how structure and management scheme affect CO₂ fluxes (Herbst et al., 2015; Musavi et al., 2017) even though it is reasonable to suppose that these are important drivers of CO₂ fluxes and that they may interact with climate and biotic variables (Luyssaert, 2014).

Here we present a case study that, in contrast, focuses on two forest sites that a) are characterized by similar site conditions, b) have a similar mean age, and c) are both dominated by beech trees (*Fagus sylvatica*) but differ in management regime and structure. We thus tackle the question of how management and related structural diversity may affect CO₂ fluxes, and directly test the hypothesis that more structurally diverse stands are less sensitive to variations in environmental and biotic drivers. This study builds on the work of Herbst et al. (2015), which was done at the same sites and showed their difference in carbon uptake and water use.

We seek to identify the major drivers of seasonal and inter-annual variability of net ecosystem productivity (NEP), gross primary productivity (GPP) and ecosystem respiration (Reco) of a structurally-diverse and a structurally-homogeneous temperate broadleaf forest. We test two hypotheses:

(1) The annual NEP and GPP of the homogeneous forest is more sensitive to variation in climate variables compared to the heterogeneous forest. A study utilizing tree rings has shown that productivity of diverse temperate beech forests exhibited higher temporal stability than monoculture forests mainly due to lower inter-annual variation as well as

due to overyielding because of asynchronous behaviour of different tree species and their interactions (Jucker et al., 2014). Grossiord et al. (2014) observed higher water availability in mixed temperate beech forests than in single species forests during drought, which they speculate as result of niche partitioning and/or facilitation processes among the interacting species.

(2) NEP and GPP of the homogeneous forest is more sensitive to intrinsic species-determined characteristics such as fruit production. Synchronous fruit production, also known as masting, is a sink for plant resources that may compete with vegetative growth (Obeso, 2002) and a negative correlation between fruit production and radial stem increment has been observed (Dittmar et al., 2003; Selås et al., 2002). Herbst et al. (2015) reported higher fruit production in the homogenous forest and here we will also quantify the effect of fruit production on annual NEP and GPP.

2. Materials and methods

1. Site description

Data were obtained from two forest sites, Hainich (DE-Hai) and Leinefelde (DE-Lnf), located in central Germany (Figure 1). The two sites are ca. 30 km apart both at an altitude of 450 mean above sea level. Soil at both sites is composed of Triassic limestone covered with variable Pleistocene loess deposits. The climate is suboceanic-submontane with a long-term annual mean air temperature of ca. 8 °C. General site characteristics are given in Table 1. The phenology of both sites is similar, with the dormant season lasting typically from November to March and growing season lasting from April to October.




Figure 1: Map showing the location of the two study sites in the central Germany. Darker patches are forests and white dots show the positions of the eddy covariance flux towers at each site. Map of Germany in inset is not to scale.

Hainich: The Hainich site (DE-Hai) is an unmanaged forest with a heterogeneous

structure, located in the central part of the Hainich National Park. Site details can be

Table 1: Instrumentation and stand characteristics for the research sites.

| Characteristics | Hainich (DE-Hai) | Leinefelde (DE-Lnf) |
|---|--|--|
| Latitude | 51°04'45.36"N | 51°19'41.58"N |
| Longitude | 10°27'07.20"E | 10°22'04.08"E |
| Altitude [m] | 440 | 450 |
| Soil | Pleistocene loess deposits with dominance of Cambisols | Pleistocene loess deposits with dominance of Luvisols |
| Instrumentation | | |
| EC measuring height [m] | 44 | 44 |
| Displacement height [m] | 22 | 22 |
| Sonic anemometer | Gill Sonic Model R3 | Gill Sonic Model R3 |
| Infra-red gas analyser (IRGA) | Li6262 | Li6262 |
| Stand characteristics | | |
| Primary species | <i>Fagus sylvatica</i> L. (64%), <i>Fraxinus excelsior</i> L. (28%), <i>Acer pseudoplatanus</i> L. (7%), and other species | <i>Fagus sylvatica</i> L. (single <i>Quercus petraea</i>) |
| Biomass [t C ha ⁻¹] | 212 | 237 |
| Plant density [trees ha ⁻¹] | 334 | 224 |
| Canopy height [m] | 35 | 35 |
| LAI [m ² m ⁻²] | 5.1 | 4.2 |
| Age (years) | Maximum up to 265, biomass weighted average = 140 | 130 ± 8 |

found in Anthoni et al. (2004) and Knohl et al. (2003). Until the end of the 19th century, it

was managed as a coppice-with-standards system and was subjected to selective cutting

until 1965. From 1965 to 1997, the area was used as a military training base and a large

part of the forest was left untouched, with only single and very valuable trees being cut.

The forest has never been clear felled and, as a result, it exhibits characteristics of an

unmanaged, old-growth forest with highly diverse horizontal and vertical structure, trees covering a wide range of age classes, up to a maximum of around 265 years, and large amounts of dead wood (both standing dead wood and coarse woody debris). The main tree species in the forest are beech (*Fagus sylvatica*, ca. 64% of tree biomass), ash (*Fraxinus excelsior*, ca. 28%), and sycamore (*Acer pseudoplatanus*, ca. 7%), with some single trees of European hornbeam (*Carpinus betulus*), elm (*Ulmus glabra*), maple (*Acer platanoides*) and other deciduous species. The main ground vegetation in the forest includes *Allium ursinum*, *Mercurialis perennis* and *Anemone nemorosa* (Mund, 2004). The Hainich flux tower site is located on a gentle north facing slope (2–3° inclination) surrounded by forest for more than 3 km in the prevailing wind direction. The only change in the surface land use is a small clearing located about 800 m perpendicular to the prevailing wind, with only 5% contribution to the overall wind direction (Knohl et al., 2003).

Leinefelde: The Leinefelde site (DE-Lnf) is an even-aged, pure beech stand managed as a shelterwood system for maximum wood production since 1938. DE-Lnf is an example of a homogenous managed even-aged 130 ± 8 years old stand that represents a late point in time of a rotation period (production cycle) lasting about 120-140 years. Crown thinning – thinning of dominant trees to reduce crowding within the main canopy – is carried out in different forest blocks regularly every 10-20 years. The last major thinning in the footprint area of the flux tower was carried out from 2002 to 2006 (Figure S1). The ground vegetation includes *Galium odoratum*, *Melica nutans*, *Milium effusum*, *Oxalis acetosella*, and *Stellaria holostea*. The eddy covariance flux tower has been in operation since April 2002. No measurements were carried out in this site from 2007 to 2009 due to access limitation.

2. Eddy covariance and meteorological measurements

Fluxes of carbon dioxide, water vapor, sensible heat, and momentum along with standard meteorological variables were measured at the two study sites. The two sites had identical eddy covariance instrumental setup and data acquisition techniques. The eddy covariance measurement system consisted of a three-dimensional sonic anemometer (Solent R3, Gill Instruments Ltd., Lymington, UK) and a fast response closed-path CO₂/H₂O infrared gas analyser in absolute mode (LI-6262, LI-COR Inc., Lincoln, NE, USA). The tube connecting the gas inlet and gas analyser was 50 m. Data were collected on a field computer using the “EddySoft” software developed by O. Kolle from MPI-Biogeochemistry, Jena, Germany (Kolle and Rebmann, 2010). Detailed information about the instrumentation can be found in Anthoni et al. (2004); Knohl et al. (2003). The turbulent fluxes were calculated using “EddyPro” software with same settings for both sites and all years. We followed Aubinet et al. (1999) and Foken et al. (2004) for quality control of the data collection and analyses. We used the “Fluxnet” online-tool (<http://www.bgc-jena.mpg.de/bgi/index.php/Services/REddyProcWeb>) and the REddyProc package in R (Wutzler et al., 2018) based on Reichstein et al. (2005) to obtain a continuous dataset of net ecosystem exchange and for partitioning this into gross primary production (GPP) and ecosystem respiration (Reco). Standard meteorological data were measured at both the sites. Anthoni et al. (2004) and Knohl et al. (2003) describe the details of the meteorological variables and instrumentation. We used the Biljou model to calculate a water availability index (WAI). Details of this model are available in Granier et al. (1999).

3. Fruit production data

Periodical fruit production is an important characteristic of beech forests. In this study, we included fruit production as a biotic variable that is assumed to affect annual carbon fluxes. The fruits (seed and pericarp) were collected in litter traps (DE-Hai 25, DE-Lnf 21

traps of 0.25 m²) distributed within the main footprint, dried at 70°C and weighed. The traps are closed plastic funnels with a small sieve at the bottom for retaining fruits while allowing drainage. The funnels are fixed on a pillar about 50 cm above the ground so that herbivores (mostly mice) are kept out.

4. Statistical analysis

Data from April 2002 to 2006 and from 2010 to 2016, the period with complete flux and meteorological data, was used to investigate the effect of meteorological variables on seasonal fluxes. We calculated zero-order correlations and slopes of the simple linear regressions between seasonal fluxes and meteorological variables. We also tested the differences of the sensitivities of seasonal CO₂ fluxes to meteorological variables between sites (Lenth and Love, 2017).

At annual scale, we used data from 2003 to 2006 and 2010 to 2016 (no fruit production data was available for 2002). Paired T-tests were applied to test for significant differences in annual CO₂ fluxes and meteorological variables between the sites. For selecting which predictor explained annual NEP, GPP and Reco, we conducted stepwise multiple linear regression (MLR) using the Akaike Information Criteria (AIC) (Field et al., 2012). The predictor variables considered for the study were mean annual air temperature (T_{air}), soil temperature (T_{soil}), global radiation (R_g), vapor pressure deficit (VPD), water availability index (WAI), spring air temperature, spring soil temperature, fruit production, growing season length, and time. The variable *time* was represented by calendar year. Thus, it integrates the effects of factors that vary and correlate with time, such as growth, effects of CO₂ fertilization (Fernández-Martínez et al., 2017), phenology (Baldocchi et al., 2018; Froelich et al., 2015; Granier et al., 2008; Pilegaard et al., 2011; Urbanski et al., 2007), and changes in nutrient deposition (Fernández-Martínez et al., 2017). To create a model applicable to both sites, we defined a MLR model using all the variables selected as significant during the per-site fits and fitted it again, this time to the combined data from


both sites. To estimate the relative effects of model predictor variables on response variables, we calculated the ‘product measure’ which distributes the overall model R^2 to each of the predictors (Nathans et al., 2012). Product measure is the product of a predictor’s β -coefficient (standardized slope) in a MLR model and its zero-order correlation. We also conducted simple regressions of predictors against the model residuals, i.e. after having removed the effects of other predictor variables via multiple linear regression. R version 3.4.3 was used for analyses (R Core Team, 2017).

3. Results

1. Meteorological characteristics


The two sites were meteorologically similar (Figure 2). Averaged yearly global radiation (R_g) from 2002 to 2016 was $122.4 \pm 6.5 \text{ W m}^{-2}$ (mean \pm sd) in DE-Hai and $124.0 \pm 7.2 \text{ W m}^{-2}$ in DE-Lnf (Figure 2a). Mean annual temperature (T_{air}) during the study period was $8.34 \pm 0.72 \text{ }^\circ\text{C}$ and $8.30 \pm 0.7 \text{ }^\circ\text{C}$ for DE-Hai and DE-Lnf, respectively. This similarity was consistent for all years with no statistically significant differences between the sites. The mean annual soil temperature (T_{soil}) was $7.61 \pm 0.36 \text{ }^\circ\text{C}$ and $8.23 \pm 0.38 \text{ }^\circ\text{C}$ for DE-Hai and DE-Lnf, respectively. A systematically higher value of $0.62 \text{ }^\circ\text{C}$ was measured in DE-Lnf ($p < 0.001$), possibly resulting from differences in the measurement depth between the sites (5 vs 4 cm in DE-Hai and DE-Lnf, respectively, Figure S2). Mean annual vapor pressure deficit (VPD) was $3.45 \pm 0.56 \text{ hPa}$ and $3.28 \pm 0.41 \text{ hPa}$ at DE-Hai and DE-Lnf, respectively, with no statistical difference between the sites. Mean annual precipitation – based on a single pluviometer per site - was significantly lower ($p < 0.001$) in DE-Lnf ($601 \pm 154 \text{ mm}$) than in DE-Hai ($744 \pm 152 \text{ mm}$). Although DE-Lnf received less rainfall, the mean annual water availability index (WAI) was similar between the sites in most years, with average values of 0.86 and 0.84 in DE-Hai and DE-

212 Lnf, respectively. We didn't observe temporal trend in any meteorological variables
 213 (Table S1).



214
 215 Figure 2: Mean annual values of (a) global radiation (R_g) and air temperature (T_{air}); (b)
 216 soil temperature (T_{soil}) and vapor pressure deficit (VPD); and (c) water availability index
 217 (WAI) and rainfall are shown from 2002 to 2016. T_{soil} was measured at 5 cm for DE-Hai
 218 and 4 cm for DE-Lnf.

219 2. Fruit production (masting) in the study sit



220
221 Figure 3: Total annual fruit production ($\text{g C m}^{-2} \text{yr}^{-1}$) in DE-Hai and DE-Lnf from 1999 to
222 2016. No data was recorded in 2001 and 2002.
223 Figure 3 shows the fruit production (masting) data for DE-Hai and DE-Lnf from 1999 to
224 2016. Data for 2001 and 2002 was not available due to technical issues. The average fruit
225 production for the entire period was $73.2 \pm 77.9 \text{ g C m}^{-2} \text{yr}^{-1}$ and $91.2 \pm 113.1 \text{ g C m}^{-2} \text{yr}^{-1}$
226 in DE-Hai and DE-Lnf, respectively. We define masting years as those when fruit
227 production is more than $50 \text{ g C m}^{-2} \text{yr}^{-1}$. Masting years occurred every two or three years.
228 Average fruit production during such years was $151 \pm 46.3 \text{ g C m}^{-2} \text{yr}^{-1}$ and $197 \pm 91.8 \text{ g}$
229 $\text{C m}^{-2} \text{yr}^{-1}$ for DE-Hai and DE-Lnf, respectively, with the former being 76 % of the latter.
230 The significant differences between DE-Lnf und DE-Hai are restricted to the
231 extraordinarily high mast years (fruit production $> 200 \text{ g C m}^{-2}$).

232 3. Cumulative net ecosystem productivity (NEP)

233 Figure 4 shows gap-filled cumulative net ecosystem productivity (NEP) data calculated
234 from eddy covariance measurements of CO_2 fluxes. Positive values correspond to a
235 cumulative net uptake of CO_2 by the vegetation (atmospheric sink) and negative values a
236 net loss (atmospheric source). At the beginning of each year, both forests are sources of
237 CO_2 . The average day of the year (doy) at which the net daily NEP switches from source

to sink (i.e. shortly after leaf-out when the rate of change in the cumulative NEP goes from negative to positive) was 125 and 121 for DE-Hai and DE-Lnf, respectively (Table S2). But this was observed as early as day 111 in DE-Hai and 112 in DE-Lnf in 2014, and as late as day 134 in DE-Hai in 2010 and day 130 in DE-Lnf in 2013. The forests continued to act as an overall C sink in average for 164 days and 168 days for DE-Hai and DE-Lnf, respectively. This growing season length was significantly correlated with mean annual temperature (Figure S3). Longer growing season length and earlier start of growing season corresponded to higher annual carbon fluxes (Figure S4c, f, i). Most of the variation in annual cumulative NEP occurred between the period of leaf out and leaf fall. The inter-annual variability in NEP during the growing season was larger in DE-Lnf compared to DE-Hai. On average, NEP in masting years was lower than in other years, with this difference being particularly visible in DE-Lnf (black solid lines in Figure 4b). The lowest cumulative NEP for both sites was measured in 2004, a year characterized by high fruit production but not the highest value observed during the study period.




Figure 4: Cumulative net ecosystem productivity for (a) DE-Hai and (b) DE-Lnf. The solid lines indicate masting years and the dashed lines indicate remaining years. The vertical grey bars indicate the time of leaf out and leaf fall. The positive values of NEP indicate a cumulative carbon uptake by the ecosystem.

4. Seasonal variability of the CO₂ fluxes

Both sites have a similar average annual cycle of NEP (Figure 5a). Each point corresponds to the mean across years of weekly sums of NEP. Both forests reached the highest carbon uptake rate between day 165 and 175. DE-Lnf showed a higher positive NEP at the start of the growing season, and DE-Hai was a stronger carbon source (black solid lines in Figure 5a) from November to December.




Figure 5: (a) The average annual NEP cycle, bar represents 95% confidence interval at 5% significance level and (b) standard deviation of weekly NEP. The grey horizontal bars show the range of the leaf out and leaf fall days of the two sites. Figure 5b shows the standard deviation of weekly NEP across years. The standard deviation of the early weeks of the year was low at both sites, and increased with the start of the growing season. A higher standard deviation in NEP was observed in DE-Lnf (dashed lines in Figure 5b) over the entire growing season. In DE-Hai (black solid lines in Figure 5b), the highest standard deviation was observed between day 215 and 230.

During the last days of the year, DE-Hai exhibited a higher standard deviation compared to DE-Lnf. For further analysis, we separated the year into different seasons, winter months (November – December: ND, January - March: JFM), spring (April – May: AM), summer (June- July: JJ), August (A) and fall (September – October: SO), to account for the different drivers and dynamics of NEP in these periods. August was treated separately from the main growing season as it is a particularly dry month and large variability observed in this period (see Figure 5b and Table 2).

5. Meteorological factors controlling seasonal variability in CO₂ fluxes

For each site, Table 2 (correlation coefficients) and Figure 6 (absolute slopes) show the effect of measured meteorological factors on CO₂ fluxes for different seasons of the year (see section 4).

Table 2: Zero-order correlation coefficients between seasonal carbon fluxes (NEP, GPP and Reco) and meteorological variables (mean of monthly values) for JFM (January - March), AM (April – May), JJ (June- July), A (August), SO (September – October) and ND (November – December) in DE-Hai and DE-Lnf. Bold numbers are statistical significant values at $p < 0.05$.

| Season | site | T _{air} | | | T _{soil} | | | R _g | | | VPD | | | WAI | | |
|--------|--------|------------------|-------------|-------------|-------------------|-------------|-------------|----------------|-------------|-------------|--------------|-------------|-------------|-------------|-------------|-------------|
| | | NEP | GPP | Reco | NEP | GPP | Reco | NEP | GPP | Reco | NEP | GPP | Reco | NEP | GPP | Reco |
| JFM | DE-Hai | -0.51 | | 0.45 | -0.52 | | 0.58 | -0.45 | | 0.34 | -0.55 | | 0.41 | | | |
| | DE-Lnf | -0.57 | | 0.78 | -0.49 | | 0.71 | -0.47 | | 0.6 | -0.66 | | 0.8 | | | |
| AM | DE-Hai | 0.85 | 0.88 | 0.82 | 0.87 | 0.91 | 0.85 | 0.6 | 0.53 | 0.34 | 0.42 | 0.33 | 0.12 | | | |
| | DE-Lnf | 0.8 | 0.85 | 0.82 | 0.84 | 0.9 | 0.87 | 0.63 | 0.59 | 0.38 | 0.37 | 0.35 | 0.23 | | | |
| JJ | DE-Hai | 0.22 | 0.38 | 0.34 | 0.27 | 0.25 | 0.1 | 0.27 | 0.51 | 0.48 | 0.11 | 0.32 | 0.36 | 0 | -0.07 | -0.1 |
| | DE-Lnf | 0.11 | 0.24 | 0.37 | -0.15 | -0.04 | 0.2 | 0.57 | 0.54 | 0.21 | 0.16 | 0.23 | 0.24 | -0.1 | 0 | 0.18 |
| A | DE-Hai | -0.36 | -0.27 | -0.02 | -0.36 | -0.24 | 0.06 | -0.38 | -0.33 | -0.15 | -0.5 | -0.49 | -0.35 | 0.65 | 0.68 | 0.54 |
| | DE-Lnf | -0.27 | -0.24 | 0.06 | -0.41 | -0.34 | 0.13 | -0.17 | -0.17 | -0.02 | -0.23 | -0.35 | -0.3 | 0.08 | 0.37 | 0.65 |
| SO | DE-Hai | 0.75 | 0.83 | 0.74 | 0.79 | 0.86 | 0.73 | 0.87 | 0.88 | 0.55 | 0.65 | 0.69 | 0.53 | -0.32 | -0.33 | -0.23 |
| | DE-Lnf | 0.62 | 0.7 | 0.81 | 0.7 | 0.77 | 0.86 | 0.76 | 0.8 | 0.75 | 0.46 | 0.52 | 0.59 | -0.19 | -0.18 | -0.12 |
| ND | DE-Hai | -0.76 | | 0.79 | -0.75 | | 0.74 | -0.78 | | 0.79 | -0.67 | | 0.76 | | | |
| | DE-Lnf | -0.69 | | 0.75 | -0.72 | | 0.76 | -0.61 | | 0.64 | -0.52 | | 0.63 | | | |

288 In winter, NEP was correlated with T_{air} , T_{soil} , R_g , and VPD at both sites with stronger
289 correlation in ND than JFM (Table 2). In ND, NEP was significantly sensitive (higher
290 absolute slopes) to T_{air} , T_{soil} and VPD in DE-Hai than in DE-Lnf (Figure 6).
291 With the arrival of spring (April – May, AM), T_{soil} became the most important factor
292 controlling spring NEP, GPP and R_{eco} in both sites, followed in importance by T_{air} , and
293 R_g . A 1 °C change in spring T_{soil} changed the NEP by 28.07 ± 6.92 and 38.74 ± 11.19 g C
294 $\text{m}^{-2} \text{month}^{-1}$ in DE-Hai and DE-Lnf, respectively. NEP was driven mainly by GPP, as
295 seen by comparing $B_{T_{\text{soil}}\&GPP}$ (slope between T_{soil} and GPP, Figure 6e) and $B_{T_{\text{soil}}\&Reco}$
296 (slope between T_{soil} and Reco, Figure 6f) for this period. NEP and GPP of DE-Lnf was
297 more sensitive to T_{soil} than DE-Hai with significantly higher $B_{T_{\text{soil}}\&NEP}$ (Figure 6d) and
298 $B_{T_{\text{soil}}\&GPP}$ (Figure 6e).
299 In summer (June – July, JJ), R_g remained a significant factor while other relationships
300 weakened. In DE-Hai, R_g showed a weak relationship with NEP but significantly
301 correlated with both GPP and Reco, while in DE-Lnf it was significantly correlated with
302 NEP and GPP. The sensitivity of NEP and GPP to R_g was higher in DE-Lnf compared to
303 DE-Hai. A change in 1 W m^{-2} of R_g increased NEP by 0.74 ± 0.47 ($p < 0.01$) g C m^{-2}
304 month^{-1} in DE-Lnf and about 0.23 ± 0.37 ($p = 0.11$) g C $\text{m}^{-2} \text{month}^{-1}$ in DE-Hai and the
305 difference between the sites is significant ($p < 0.1$).
306 In August (A), only the soil water availability index (WAI) correlated significantly with
307 CO₂ fluxes, specifically with NEP, GPP and R_{eco} in DE-Hai and only with Reco in DE-
308 Lnf.
309 In fall (September – October, SO), T_{air} , T_{soil} , R_g and VPD, again became significantly
310 correlated with NEP, GPP and R_{eco} in both sites, with high correlation values for the first
311 three and slightly lower ones for VPD.




Figure 6: Slopes (B) between seasonal CO₂ fluxes (NEP – column 1, GPP – column 2 and Reco – column 3) and meteorological variables (mean of monthly values) for JFM (January - March), AM (April – May), JJ (June- July), A (August), SO (September – October) and ND (November – December). The larger circles and squares indicate slopes, which are significantly different from zero. *** indicate statistical significant difference of slopes between two sites at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$ at bottom of each figure (if any). The first letter of each month is used in the timescale.

6. Seasonal differences in CO₂ fluxes between the two sites




Figure 7: Difference in sum of (a) NEP, (b) GPP and (c) Reco between the two sites for JFM (January - March), AM (April – May), JJ (June- July), A (August), SO (September – October) and ND (November – December). The Y-axis represents $\Delta X = X_{\text{DE-Lnf}} - X_{\text{DE-Hai}}$ (X being seasonal NEP, GPP or Reco). Bars represent 95% confidence interval. The dashed line marks the zero line (indicates no difference between the sites). *** indicate statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.

In addition to slopes, we also looked at differences of average seasonal sums of NEP,

GPP and Reco between sites (Figure 7). Significantly higher NEPs observed in DE-Lnf

than in DE-Hai in JFM and ND (5.48 and $17.79 \text{ g C m}^{-2} \text{ month}^{-1}$, Figure 7a) was due to

lower Reco in DE-Lnf (Figure 7c). DE-Lnf also acted as a stronger carbon sink in the

spring ($26 \text{ g C m}^{-2} \text{ month}^{-1}$) which was mostly due to higher GPP of $24 \text{ g C m}^{-2} \text{ month}^{-1}$.

In JJ, difference between NEP in two sites were insignificant because both GPP and Reco

in DE-Lnf were significantly higher by $19.08 \text{ g C m}^{-2} \text{ month}^{-1}$ and $10.94 \text{ g C m}^{-2} \text{ month}^{-1}$,

respectively. During August, NEP and GPP were similar in both sites, however, a higher Reco was observed in DE-Lnf ($10.04 \text{ g C m}^{-2} \text{ month}^{-1}$). In the fall, NEP, GPP and Reco were lower in DE-Lnf but differences were insignificant.

7. Annual estimates of NEP, GPP and Reco

Annual gap-filled NEP, GPP and Reco for both sites are presented in Figure 8. Mean Annual NEP was 487 ± 57.8 (mean \pm SD for the mean of all years) and $585 \pm 144 \text{ g C m}^{-2} \text{ yr}^{-1}$ in DE-Hai and DE-Lnf, respectively. A paired t-test showed that DE-Lnf was a significantly stronger carbon sink ($98 \text{ g C m}^{-2} \text{ yr}^{-1}$, $p < 0.05$). We observed the largest differences in NEP between the two sites from 2010 (Table S3).

Annual NEP was significantly more variable in DE-Lnf (Levene's test with $p < 0.05$), with coefficients of variation (CV) being 12% and 25 % in DE-Hai and DE-Lnf, respectively. A significant temporal trend of NEP was observed only for DE-Lnf, with an increase of $21.8 \text{ g C m}^{-2} \text{ yr}^{-1}$ ($p < 0.05$).

The mean of annual GPP values at DE-Hai and DE-Lnf was 1559 ± 118 and $1627 \pm 164 \text{ g C m}^{-2} \text{ yr}^{-1}$ respectively, with the difference being statistically significant ($p = 0.07$). Like NEP, significantly higher annual GPP was again observed in DE-Lnf from 2010 (Table S3). The variability in annual GPP was not significantly different between the two sites, with coefficients of variation (CV) equalling 7.5 % and 10 % in DE-Hai and DE-Lnf, respectively. As for annual NEP, no temporal trend in annual GPP was observed in DE-Hai, whereas a significant trend of $25.5 \text{ g C m}^{-2} \text{ yr}^{-1}$ ($p < 0.05$) was observed in DE-Lnf.

The mean annual ecosystem respiration (Reco) was 1071 ± 96 and $1042 \pm 60 \text{ g C m}^{-2} \text{ yr}^{-1}$ in DE-Hai and DE-Lnf, respectively with no significant difference between sites. In contrast to annual NEP and GPP, annual Reco was lower in DE-Lnf with no significant difference. Annual Reco showed a positive temporal trend at both sites but it was statistically insignificant (Table S1).




Figure 8: Annual sums of GPP, Reco and NEP from 2003 to 2016. Data from 2007 to 2009 was not measured in DE-Lnf.

8. Factors contributing to annual variability of NEP, GPP and Reco

Together, fruit production (FP), time (see the section 2.4) and mean annual soil temperature (T_{soil}) explained ca. 65% and 92% of the variation in annual NEP in DE-Hai and DE-Lnf, respectively (Table 3). In DE-Hai, T_{soil} was the most important factor, followed by FP (negative correlation). In the case of DE-Lnf, time and FP explained most of the variation.

Results were similar for annual GPP, with a total R^2 of 0.62 and 0.88 in DE-Hai and DE-Lnf, respectively. T_{soil} was the only significant variable for DE-Hai, explaining most of the variation in GPP. In DE-Lnf, time was the strongest predictor variable, followed by T_{soil} and FP (negative correlation). A significant relationship between annual Reco was found only for DE-Lnf with T_{soil} as the only significant variable.

Figure 9 shows the relationship between each of the three significant driving variables (T_{soil} , FP, time) and the three flux quantities (NEP, GPP, Reco) in terms of the residual variance remaining after the effects of the other two driving variables (e.g. FP and time, in the case of T_{soil}) have been removed. This analysis increased the amount of variation in NEP and GPP explained by T_{soil} . We tested the difference between the slopes of two sites obtained in Figure 9 (Table S4). The sensitivities of NEP on T_{soil} were similar between

the sites when the effect of fruit production and time is removed. The result was similar for GPP. Similarly, fruit production was significantly correlated with residuals of NEP after removing effect of T_{soil} and time (Figure 9b) in both sites. We observed that residuals of GPP decreased with increased fruit production, but significant only for DE-Lnf and with a more than twice as large slope (Figure 9e and Table S4). Slope between residuals of annual Reco and fruit production was not significantly different from zero for both sites but had a positive slope for DE-Lnf. Only in DE-Lnf, we observed positive slope between time and residuals of NEP after removing the effect of T_{soil} and FP (Figure 9c and Table S4). Higher slope of residuals of

Table 3: Major factors contributing to the variation of annual NEP, GPP and Reco. Var is the contribution of each predictor to total R^2 calculated with the product measure metric.

| Flux | Predictors | DE-Hai | | | DE-Lnf | | |
|------|---------------------|--------|--------------|------|--------|--------------|------|
| | | R^2 | Coefficients | Var | R^2 | Coefficients | Var |
| NEP | | 0.65 | | | 0.92 | | |
| | T_{soil} + | | 110.48 * | 0.49 | | 79.94 (*) | 0.07 |
| | FP + | | -0.34 (*) | 0.15 | | -0.74 *** | 0.34 |
| | Time | | -0.99 | 0.00 | | 22.30 *** | 0.51 |
| GPP | | 0.62 | | | 0.88 | | |
| | T_{soil} + | | 208.08 * | 0.49 | | 189.95 * | 0.28 |
| | FP + | | -0.23 | 0.00 | | -0.60 ** | 0.13 |
| | Time | | 8.93 | 0.13 | | 22.57 * | 0.46 |
| Reco | | 0.45 | | | 0.79 | | |
| | T_{soil} + | | 97.61 | 0.19 | | 109.90 ** | 0.65 |
| | FP + | | 0.02 | 0.00 | | 0.14 | 0.14 |
| | Time | | 9.92 | 0.25 | | 0.27 | 0.00 |

*** significant at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; (*) significant at $p < 0.1$

GPP (Figure 9f) compared to residuals of Reco (Figure 9i) to that. In DE-Hai, positive slope between time and residuals of GPP was negated by Reco, thus showing no effect in residuals of NEP.




Figure 9: Simple linear regressions between residuals of different carbon fluxes after removing effect of fruit production and time (1st column, a, d, g), T_{soil} and time (2nd column, b, e, h) or T_{soil} and fruit production (3rd column, c, f, i). Here time is represented by number plus 2000 for clean graph. The solid lines represent linear regression lines and dashed lines confidence interval at 5% significance level. *** indicate statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.

1. Discussion

1. Meteorological controls of seasonal CO₂ fluxes

We observed stronger sensitivity of winter Reco to T_{soil} (Figure 6) resulting in higher Reco in DE-Hai than in DE-Lnf, thus, higher winter NEP in DE-Lnf (Figure 7). Higher

ecosystem respiration is most likely driven by higher stocks of decaying dead wood and by higher rates of leaf litter decomposition caused by higher leaf litter quality (proportion of litter from ash and maple trees) and a higher biological activity in the soil (Mund, 2004).

In spring, NEP, GPP and Reco strongly correlated with T_{air} and T_{soil} with higher correlation values for T_{soil} in both sites. The change of NEP with T_{soil} was mostly driven by the sensitivity of GPP to T_{soil} . Similar results were also observed in Borden forest, Canada, during the spring time (Froelich et al., 2015). Spring temperature is very important for leaf unfolding and we found that the growing season started earlier when spring air temperatures were higher (Figure S5). Also, warm soil means favourable growth conditions for roots (Alvarez-Uria and Körner, 2007), improving nutrient and water uptake and leading to increased photosynthesis. Along with GPP, Reco also increases with increased soil temperature, due to the increased activities of tree roots and microbes (Davidson et al., 1998; Gonzalez-Meler et al., 2013). DE-Lnf was a significantly stronger sink during spring, primarily driven by GPP (Figure 7). We further observed higher sensitivity of GPP to T_{soil} in DE-Lnf than in DE-Hai ($p < 0.1$). This could be explained by an earlier physiological activity in beech than in ash (Cole and Sheldon, 2017). DE-Lnf is a beech monoculture comprising mostly vital trees at their optimal age (optimal regarding wood growth and fruit production) whereas DE-Hai has 28% ash trees and includes many small, young, suppressed, very old, semi-dead and dead trees.

In summer (June - July), R_g was the most important environmental factor controlling carbon fluxes of both sites. DE-Lnf was a stronger sink with higher GPP and Reco because it is full of optimally growing trees that may have capitalised the radiation more than DE-Hai. Similarly, the stronger sensitivity of summer GPP and Reco to R_g of DE-

Lnf can be attributed to optimally growing trees of similar size and age that react to weather conditions in same direction and magnitude. August is of interest for DE-Hai where soil water availability (WAI) influences its CO₂ fluxes. DE-Hai is more affected than DE-Lnf by water availability because of ash trees that are still active when WAI drops below a critical value (e.g. 2003). Trees close their stomata as the soil water availability reduces to prevent water loss (Chaves et al., 2002) also decreasing photosynthesis and respiration.

2. Annual CO₂ fluxes and its inter-annual variability

Before comparing DE-Lnf with DE-Hai, it is useful to discuss site management and history. DE-Lnf is an example of a managed even-aged stand that represents one point in time of a rotation period (production cycle) lasting about 120-140 years. Within the production cycle, DE-Lnf represents a mature stand at its late optimum phase - with “late optimum” meaning close to the end of highest timber production. If the stand were not a certified seed production site, and if the eddy tower were not there, the stand would have been thinned more heavily (preparatory thinning) to start the regeneration under the shelter of the older trees. This means that if the site were like any other common managed beech stand, then living biomass and NEP might be much smaller. For a comparison of managed, even-aged forests with unmanaged forests a chronosequence of several even-aged stands covering the entire production cycle would be needed. We thus note that the heterogeneous stand is not being compared with the average managed beech forest in the region, but rather with a particularly mature and productive stand. This must be taken into account before concluding on the general impact of management on the C cycle.

Both the sites were strong carbon sinks despite one site being in an advanced stage of growth with trees as old as 265 years. The ranges of annual NEP were comparable with Oak Ridge forest in Tennessee, US ($577 \pm 63 \text{ g C m}^{-2} \text{ yr}^{-1}$, Wilson and Baldocchi, 2001), Ozarks forest in Missouri, US ($479 \pm 65 \text{ g C m}^{-2} \text{ yr}^{-1}$, Shao et al., 2014), and Oak

460 woodland forests, UK ($486 \pm 115 \text{ g C m}^{-2} \text{ yr}^{-1}$, Wilkinson et al., 2012) but slightly higher
461 than average annual NEP for temperate forests ($350 \pm 100 \text{ g C m}^{-2} \text{ yr}^{-1}$, Table S5).
462 Average annual GPP and Reco of both sites also fell within the range average annual
463 GPP ($1506 \pm 214 \text{ g C m}^{-2} \text{ yr}^{-1}$) and Reco ($1181 \pm 158 \text{ g C m}^{-2} \text{ yr}^{-1}$) of temperate
464 deciduous forests.

465 A previous study of the same sites using seven years of data (Herbst et al., 2015) reported
466 small but non-significant difference between the sites for NEP. However, by including
467 four more recent years we found a significant difference in NEP between the two sites.
468 The absolute difference in mean annual NEP ($98 \text{ g C m}^{-2} \text{ yr}^{-1}$) results from a higher mean
469 annual GPP of $69 \text{ g C m}^{-2} \text{ yr}^{-1}$ (2/3 of NEP) plus a lower mean annual Reco of 29 g C m^{-2}
470 yr^{-1} (1/3 of NEP) in DE-Lnf. The higher mean annual values of NEP in DE-Lnf is due to
471 higher annual NEP after 2011 than DE-Hai. After 2011, significantly higher GPP and
472 lower Reco in DE-Lnf resulted in higher NEP. The higher carbon uptake in DE-Lnf was
473 determined by the activities during winter, spring and summer (section 1).

474 We observed lower coefficient of variation (CV) of annual NEP for both the sites
475 compared to average CV of annual NEP of temperate forests. Average CV of annual NEP
476 for temperate forests is 35 % ($\text{sd} = \pm 100 \text{ g C m}^{-2} \text{ yr}^{-1}$, Baldocchi et al., 2018 and Table
477 S5) with the highest CV of 66 % observed in the Borden forest (Froelich et al., 2015) and
478 Sorø forest (Pilegaard et al., 2011). Interestingly, the CV of annual NEP in DE-Hai was
479 the lowest among reported results for temperate deciduous forests and remained similar
480 even after adding four years of data. Also, the CV of annual GPP and Reco of both sites
481 were lower than the average for temperate deciduous forests. Like annual NEP, CV of
482 annual GPP in DE-Hai was lowest among all the temperate deciduous broadleaved
483 forests following the results of Musavi et al. (2017) that reported older and diverse forests
484 had less variation in saturated gross primary productivity (GPP_{sat}). Between our sites, the

managed, homogeneous forest showed a higher CV in NEP and GPP than the unmanaged, heterogeneous forest.

Long-term studies conducted in temperate deciduous forests have identified many factors contributing to site-specific inter-annual variation of NEP. Some studies have found that growing season length explains inter-annual variation in NEP: the Borden forest in Canada (Froelich et al., 2015), Sorø forest in Denmark (Pilegaard et al., 2011), Hesse forest (Granier et al., 2008), Morgan-Monroe State Forest in Indiana (Dragoni et al., 2011). In our case, we observed positive correlation between growing season length and NEP and GPP (Figure S4e, f and g), and at the same time positive correlation between growing season length and temperatures (Figure S3). Among T_{air} and T_{soil} , we found stronger relationship of T_{soil} with NEP and GPP (Figure S6 and Figure S7), this could have led to selection of T_{soil} in multiple linear model selection using AIC criteria. At an annual scale, we found that sensitivities of T_{soil} to NEP and GPP were similar based on residual analysis conducted after removing effect of fruit production and time. Other two important factors that explained annual fluxes were fruit production and time. We will discuss them separately in section 3 and 4.

3. Effect of fruit production on CO_2 fluxes

Fruit production (FP) was negatively correlated with NEP at both sites (Figure S8). The negative slope of NEP vs fruit production was mostly the result of a reduction of GPP in high fruit production years, i.e. photosynthesis decreased with increasing fruit production. Many studies reported that tree ring growth was reduced in years of high fruit production (e.g. Holmsgaard, 1955; Mund et al., 2010). Different mechanisms have been reported for this reduced growth in trees, including reduced photosynthetic rates in reproductive branches due to N or P depletion in those branches (Sala et al., 2012), smaller leaves (Innes, 1992), reduced leaf area (Ferretti et al., 1998), lower number of leaves during the

most year (Han et al., 2008) which could be due to diminished shoot growth and increased foliar bud mortality (Ishihara and Kikuzawa, 2009) as foliar buds are replaced by seeds (Innes, 1994). It is important to note that the effect was stronger in DE-Lnf - a homogeneous forests with trees in a similar fruit-producing age class (150 -170 years) (Herbst et al., 2015). On the other hand, DE-Hai has a heterogeneous structure with different species and a wide range of tree age from 0 to 265 years. Thus, fruit production does not occur in all trees (Figure 3).

4. Effect of time on CO₂ fluxes

Many studies report that CO₂ fluxes in temperate forests have been increasing (Fernández-Martínez et al., 2017; Froelich et al., 2015; Granier et al., 2008; Pilegaard et al., 2011b, etc). A significant increasing temporal trend in CO₂ uptake has also been observed in ca. 80 year-old managed beech forest in Sorø, Denmark (-23 g C m⁻² yr⁻¹, Pilegaard et al., 2011), ca. 40 year-old managed beech forest in Hesse, France (-43 g C m⁻² yr⁻¹, Granier et al., 2008) and ca. 95 year-old managed maple, white oak and red oak Harvard forest in US (-16 g C m⁻² yr⁻¹, Urbanski et al., 2007). We observed a similar temporal trend in NEP in DE-Lnf, a managed homogeneous forest, which was not visible when Herbst et al., (2015) reported 7 years' data. The NEP of DE-Hai was reported to be - 494 g C m⁻² and - 490 g C m⁻² for 2000 and 2001, respectively by Knohl et al., (2003) and the capacity remained in the same range for all 12 years described here, exhibiting no significant temporal trend. However, we observed positive trends in annual GPP and Reco at DE-Hai which might have cancelled resulting in no trend in NEP. The normal temporal trend in forest productivity is to follow a sigmoidal growth curve as individuals age. Overlaid on this, however, are the impacts of increasing atmospheric CO₂ concentration, decreasing sulphur deposition (Fernández-Martínez et al., 2017), increasing nitrogen deposition, as well as management activities such as thinning. Here

we used time as a variable because it is hard to disentangle these effects. Thus, our observed temporal trend in DE-Lnf needs a careful interpretation because the trend appears to reflect an increase in CO₂ uptake starting from 2012 (Figure 8), and there was no significant temporal trend in any of the observed meteorological variables that could explain this increase (Table S1). We observed a positive trend in growing season length (0.75 days per year, $p < 0.05$), which explained about 19% of the variability in NEP ($p = 0.16$) but which was not as high as reported for Hesse (Granier et al., 2008), Sorø (Pilegaard et al., 2011), Borden forest (Froelich et al., 2015), and Morgan Monroe state forest (Dragoni et al., 2011). As an alternate hypothesis, we note that thinning operations were carried out in the main flux footprint area of DE-Lnf, and we speculate that because thinning has the effect of increasing productivity in the remaining trees, this might have impacted NEP subsequently. About 2319 g C m⁻² of biomass was thinned from the footprint area from 2002 to 2006 (Figure S1) and the largest thinning was carried out in 2005 (998 g C m⁻²). We observed higher NEP of 171 g C m⁻² yr⁻¹ in the period 2010 – 2016 compared to 2003 – 2006 (Table S3). About 1197 g C m⁻² of additional carbon was absorbed by DE-Lnf over the period of 7 years. We note that Sorø was thinned about 20 % every 10 years (Pilegaard et al., 2011) and Hesse was thinned every five years (Granier et al., 2008), yet these authors did not report any significant effect of thinning on CO₂ fluxes. This could be due to the effects of thinning being seen only gradually over the following years as trees adjust to the new conditions, thus making the connection between C fluxes and thinning difficult to perceive or quantify.

2. Conclusions

We compared two temperate deciduous forest types with similar site and meteorological conditions but with different structure in terms of diameter distribution, age and species composition. We found that the homogeneous forest was a higher carbon sink than the heterogeneous forest due to lower respiration rates in winter and higher carbon uptake

rates in spring and summer. CO₂ uptake by the homogeneous forest has increased in recent years playing a key role in determining differences between the forests. We identified an overall time-effect but could not disentangle possible contributing factors such as increasing atmospheric CO₂ concentrations or effects of silvicultural management.

In the introduction section, we put forward two hypotheses. Regarding hypothesis one, i.e. higher sensitivity of carbon fluxes of homogenous forests to environmental variables, we concluded that the homogenous forest showed a stronger sensitivity to environmental variables during spring (T_{soil}) and summer (R_g) causing inter-annual differences between sites. At annual scale, however, the sensitivities of CO₂ fluxes to environment variables are similar due to stronger control by biotic factors. In case of the second hypothesis, i.e. a higher negative sensitivity of CO₂ fluxes of the homogenous forest to fruit production, we see that the NEP of the homogenous forest showed a stronger sensitivity to fruit production due to a higher negative sensitivity of GPP and higher positive sensitivity of Reco to fruit production. The relationships are weak; thus, we suggest that more data are required to confirm the hypothesis.

Even though both forests are of same average age, structures of these forests vary. This leads to different responses of the CO₂ fluxes to environmental and biotic factors. Thus, it is necessary that we include structural information along with species traits (fruiting characteristics) and management activities to be able to predict the CO₂ fluxes in response to future climate.

Acknowledgements

We would like to thank Erasmus Mundus Joint Doctorate Programme Forest and Nature for Society (EMJD FONASO) and German Federal Ministry of Education and Research for funding this research. We thank the administration of the Hainich National Park and the forestry district Leinefelde for the opportunity for research in their forest areas. We

588 are also indebted to Ernst-Detlef Schulze, Olaf Kolle, Kerstin Hippler, Karl Kübler,
589 Martin Hertel, Agnes Fastnacht (Max-Planck Institute for Biogeochemistry), Peter
590 Anthoni (Karlsruhe Institute of Technology KIT, Institute of Meteorology and Climate
591 Research Atmospheric Environmental Research), Corinna Rebmann (Helmholtz Centre
592 for Environmental Research – UFZ), Frank Tiedemann, Dietmar Fellert, Heinrich
593 Kreilein, Martin Lindenberg, Lukas Siebicke (University of Göttingen) and Werner
594 Kutsch (ICOS) for their work at the Hainich and Leinefelde tower sites.

595

- 597 Alvarez-Uria, P., Körner, C., 2007. Low temperature limits of root growth in deciduous
598 and evergreen temperate tree species. *Funct. Ecol.* 21, 211–218.
599 <https://doi.org/10.1111/j.1365-2435.2007.01231.x>
- 600 Anthoni, P.M., Knohl, A., Rebmann, C., Freibauer, A., Mund, M., Ziegler, W., Kolle, O.,
601 Schulze, E.D., 2004. Forest and agricultural land-use-dependent CO₂ exchange in
602 Thuringia, Germany. *Glob. Chang. Biol.* 10, 2005–2019.
603 <https://doi.org/10.1111/j.1365-2486.2004.00863.x>
- 604 Aubinet, M., Grelle, A., Ibrom, A., Rannik, Ü., Moncrieff, J., Foken, T., Kowalski, A.S.,
605 Martin, P.H., Berbigier, P., Bernhofer, C., Clement, R., Elbers, J., Granier, A.,
606 Grünwald, T., Morgenstern, K., Pilegaard, K., Rebmann, C., Snijders, W., Valentini,
607 R., Vesala, T., 1999. Estimates of the annual net carbon and water exchange of
608 forests: the EUROFLUX methodology. *Adv. Ecol. Res.* 30, 113–175.
609 [https://doi.org/10.1016/S0065-2504\(08\)60018-5](https://doi.org/10.1016/S0065-2504(08)60018-5)
- 610 Baldocchi, D., Chu, H., Reichstein, M., 2018. Inter-annual variability of net and gross
611 ecosystem carbon fluxes: A review. *Agric. For. Meteorol.* 249, 520–533.
612 <https://doi.org/10.1016/j.agrformet.2017.05.015>
- 613 Baldocchi, D., Falge, E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P.,
614 Bernhofer, C., Davis, K., Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee,
615 X., Malhi, Y., Meyers, T., Munger, W., Oechel, W., Paw, K.T., Pilegaard, K., Schmid,
616 H.P., Valentini, R., Verma, S., Vesala, T., Wilson, K., Wofsy, S., Baldocchi, D., Falge,
617 E., Gu, L., Olson, R., Hollinger, D., Running, S., Anthoni, P., Bernhofer, C., Davis, K.,
618 Evans, R., Fuentes, J., Goldstein, A., Katul, G., Law, B., Lee, X., Malhi, Y., Meyers, T.,
619 Munger, W., Oechel, W., Paw, K.T., Pilegaard, K., Schmid, H.P., Valentini, R., Verma,
620 S., Vesala, T., Wilson, K., Wofsy, S., 2001. FLUXNET: A new tool to study the
621 temporal and spatial variability of ecosystem–scale carbon dioxide, water vapor,
622 and energy flux densities. *Bull. Am. Meteorol. Soc.* 82, 2415–2434.
623 [https://doi.org/10.1175/1520-0477\(2001\)082<2415:FANTTS>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<2415:FANTTS>2.3.CO;2)
- 624 Baldocchi, D., Xu, L., 2005. Carbon exchange of deciduous broadleaved forestes in
625 temperate and Mediterranean regions, in: Griffiths, H., Jarvis, P. (Eds.), *The Carbon*
626 *Balance of Forest Biomes*. Taylor Francis, Trowbridge, UK, pp. 187–214.
- 627 Barr, A.G., Black, T.A., Hogg, E.H., Griffis, T.J., Morgenstern, K., Kljun, N., Theede, A.,
628 Nesic, Z., 2007. Climatic controls on the carbon and water balances of a boreal
629 aspen forest, 1994–2003. *Glob. Chang. Biol.* 13, 561–576.
630 <https://doi.org/10.1111/j.1365-2486.2006.01220.x>
- 631 Beer, C., Reichstein, M., Tomelleri, E., Ciais, P., Jung, M., Carvalhais, N., Rodenbeck, C.,
632 Arain, M.A., Baldocchi, D., Bonan, G.B., Bondeau, A., Cescatti, A., Lasslop, G.,
633 Lindroth, A., Lomas, M., Luyssaert, S., Margolis, H., Oleson, K.W., Rouspard, O.,
634 Veenendaal, E., Viovy, N., Williams, C., Woodward, F.I., Papale, D., 2010. Terrestrial
635 Gross Carbon Dioxide Uptake: Global Distribution and Covariation with Climate.
636 *Science* (80-.). 329, 834–838. <https://doi.org/10.1126/science.1184984>
- 637 Chaves, M.M., Pereira, J.S., Maroco, J., Rodrigues, M.L., Ricardo, C.P.P., Osório, M.L.,
638 Carvalho, I., Faria, T., Pinheiro, C., 2002. How plants cope with water stress in the
639 field. *Photosynthesis and growth*. *Ann. Bot.* 89, 907–916.
640 <https://doi.org/10.1093/aob/mcf105>
- 641 Chen, Z., Yu, G., Zhu, X., Wang, Q., Niu, S., Hu, Z., 2015. Covariation between gross
642 primary production and ecosystem respiration across space and the underlying

mechanisms: A global synthesis. *Agric. For. Meteorol.* 203, 180–190.
<https://doi.org/10.1016/j.agrformet.2015.01.012>

Chu, H., Chen, J., Gottgens, J.F., Desai, A.R., Ouyang, Z., Qian, S.S., 2016. Response and biophysical regulation of carbon dioxide fluxes to climate variability and anomaly in contrasting ecosystems in northwestern Ohio, USA. *Agric. For. Meteorol.* 220, 50–68. <https://doi.org/10.1016/j.agrformet.2016.01.008>

Ciais, P., Reichstein, M., Viovy, N., Granier, A., Ogée, J., Allard, V., Aubinet, M., Buchmann, N., Bernhofer, C., Carrara, A., Chevallier, F., De Noblet, N., Friend, A.D., Friedlingstein, P., Grünwald, T., Heinesch, B., Keronen, P., Knohl, A., Krinner, G., Loustau, D., Manca, G., Matteucci, G., Miglietta, F., Ourcival, J.M., Papale, D., Pilegaard, K., Rambal, S., Seufert, G., Soussana, J.F., Sanz, M.J., Schulze, E.D., Vesala, T., Valentini, R., 2005. Europe-wide reduction in primary productivity caused by the heat and drought in 2003. *Nature* 437, 529–533.
<https://doi.org/10.1038/nature03972>

Cole, E.F., Sheldon, B.C., 2017. The shifting phenological landscape: Within- and between-species variation in leaf emergence in a mixed-deciduous woodland. *Ecol. Evol.* 7, 1135–1147. <https://doi.org/10.1002/ece3.2718>

Davidson, E.A., Belk, E., Boone, R.D., 1998. Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest. *Glob. Chang. Biol.* 4, 217–227.
<https://doi.org/10.1046/j.1365-2486.1998.00128.x>

Dittmar, C., Zech, W., Elling, W., 2003. Growth variations of Common beech (*Fagus sylvatica* L.) under different climatic and environmental conditions in Europe - A dendroecological study. *For. Ecol. Manage.* 173, 63–78.
[https://doi.org/10.1016/S0378-1127\(01\)00816-7](https://doi.org/10.1016/S0378-1127(01)00816-7)

Dragoni, D., Schmid, H.P., Wayson, C.A., Potter, H., Grimmond, C.S.B., Randolph, J.C., 2011. Evidence of increased net ecosystem productivity associated with a longer vegetated season in a deciduous forest in south-central Indiana, USA. *Glob. Chang. Biol.* 17, 886–897. <https://doi.org/10.1111/j.1365-2486.2010.02281.x>

Fernández-Martínez, M., Vicca, S., Janssens, I.A., Ciais, P., Obersteiner, M., Bartrons, M., Sardans, J., Verger, A., Canadell, J.G., Chevallier, F., Wang, X., Bernhofer, C., Curtis, P.S., Gianelle, D., Grünwald, T., Heinesch, B., Ibrom, A., Knohl, A., Laurila, T., Law, B.E., Limousin, J.M., Longdoz, B., Loustau, D., Mammarella, I., Matteucci, G., Monson, R.K., Montagnani, L., Moors, E.J., Munger, J.W., Papale, D., Piao, S.L., Peñuelas, J., 2017. Atmospheric deposition, CO₂, and change in the land carbon sink. *Sci. Rep.* 7, 1–13. <https://doi.org/10.1038/s41598-017-08755-8>

Ferretti, M., Baratozzi, L., Cenni, E., Cozzi, A., Savini, P., 1998. Crown transparency of beech (*Fagus sylvatica* L.) in the northern Apennines (Italy) - Status, changes and relationships with site characteristics and other indices of tree condition. *Chemosphere* 36, 1037–1042. [https://doi.org/10.1016/S0045-6535\(97\)10168-0](https://doi.org/10.1016/S0045-6535(97)10168-0)

Field, A.P., Miles, J., Field, Z., 2012. *Discovering statistics using R*. Sage.

Foken, T., Göckede, M., Mauder, M., Mahrt, L., Amiro, B., Munger, W., 2004. Post-field data quality control, in: *Handbook of Micrometeorology*. Kluwer Academic Publishers, Dordrecht, pp. 181–208. https://doi.org/10.1007/1-4020-2265-4_9

Froelich, N., Croft, H., Chen, J.M., Gonsamo, A., Staebler, R.M., 2015. Trends of carbon fluxes and climate over a mixed temperate-boreal transition forest in southern Ontario, Canada. *Agric. For. Meteorol.* 211–212, 72–84.
<https://doi.org/10.1016/j.agrformet.2015.05.009>

691 Gonzalez-Meler, M.A., Hopkins, F., Flower, C.E., Lynch, D.J., Czimczik, C., Tang, J., Subke,
692 J.-A., 2013. Ecosystem-level controls on root-rhizosphere respiration. *New Phytol.*
693 199, 339–351. <https://doi.org/10.1111/nph.12271>

694 Granier, A., Bréda, N., Biron, P., Villetle, S., 1999. A lumped water balance model to
695 evaluate duration and intensity of drought constraints in forest stands. *Ecol.*
696 *Modell.* 116, 269–283. [https://doi.org/10.1016/S0304-3800\(98\)00205-1](https://doi.org/10.1016/S0304-3800(98)00205-1)

697 Granier, A., Bréda, N., Longdoz, B., Gross, P., Ngao, J., 2008. Ten years of fluxes and
698 stand growth in a young beech forest at Hesse, North-eastern France. *Ann. For. Sci.*
699 65, 704. <https://doi.org/10.1051/Forest:2008052>

700 Grossiord, C., Granier, A., Ratcliffe, S., Bouriaud, O., Bruelheide, H., Checko, E., Forrester,
701 D.I., Dawud, S.M., Finer, L., Pollastrini, M., Scherer-Lorenzen, M., Valladares, F.,
702 Bonal, D., Gessler, A., 2014. Tree diversity does not always improve resistance of
703 forest ecosystems to drought. *Proc. Natl. Acad. Sci.* 111, 14812–14815.
704 <https://doi.org/10.1073/pnas.1411970111>

705 Han, Q., Kabeya, D., Iio, A., Kakubari, Y., 2008. Masting in *Fagus crenata* and its influence
706 on the nitrogen content and dry mass of winter buds. *Tree Physiol.* 28, 1269–1276.
707 <https://doi.org/10.1093/treephys/28.8.1269>

708 Herbst, M., Mund, M., Tamrakar, R., Knohl, A., 2015. Differences in carbon uptake and
709 water use between a managed and an unmanaged beech forest in central
710 Germany. *For. Ecol. Manage.* 355, 101–108.
711 <https://doi.org/10.1016/j.foreco.2015.05.034>

712 Holmsgaard, E., 1955. Tree-ring analyses of Danish forest trees. *Det Forstl. forsøgsvæsen*
713 *i Danmark XXII*, 1–246.

714 Hommeltenberg, J., Schmid, H.P., Drösler, M., Werle, P., 2014. Can a bog drained for
715 forestry be a stronger carbon sink than a natural bog forest? *Biogeosciences* 11,
716 3477–3493. <https://doi.org/10.5194/bg-11-3477-2014>

717 Hui, D., Luo, Y., Katul, G., 2003. Partitioning interannual variability in net ecosystem
718 exchange between climatic variability and functional change 433–442.

719 Humphreys, E.R., Lafleur, P.M., 2011. Does earlier snowmelt lead to greater CO₂
720 sequestration in two low Arctic tundra ecosystems? *Geophys. Res. Lett.* 38, n/a-
721 n/a. <https://doi.org/10.1029/2011GL047339>

722 IGBP Terrestrial Carbon Working Group, I.T.C.W., 1998. CLIMATE: The Terrestrial Carbon
723 Cycle: Implications for the Kyoto Protocol. *Science* (80-.). 280, 1393–1394.
724 <https://doi.org/10.1126/science.280.5368.1393>

725 Innes, J.L., 1994. The occurrence of flowering and fruiting on individual trees over 3
726 years and their effects on subsequent crown condition. *Trees* 8, 139–150.
727 <https://doi.org/10.1007/BF00196638>

728 Innes, J.L., 1992. Observations on the condition of beech (*Fagus sylvatica* L.) in Britain in
729 1990. *Forestry* 65, 35–60. <https://doi.org/10.1093/forestry/65.1.35>

730 Ishihara, M.I., Kikuzawa, K., 2009. Annual and spatial variation in shoot demography
731 associated with masting in *Betula grossa*: Comparison between mature trees and
732 saplings. *Ann. Bot.* 104, 1195–1205. <https://doi.org/10.1093/aob/mcp217>

733 Jensen, R., Herbst, M., Friborg, T., 2017. Direct and indirect controls of the interannual
734 variability in atmospheric CO₂ exchange of three contrasting ecosystems in
735 Denmark. *Agric. For. Meteorol.* 233, 12–31.
736 <https://doi.org/10.1016/j.agrformet.2016.10.023>

737 Jucker, T., Bouriaud, O., Avacaritei, D., Coomes, D.A., 2014. Stabilizing effects of diversity
738 on aboveground wood production in forest ecosystems: linking patterns and

739 processes. *Ecol. Lett.* 17, 1560–1569. <https://doi.org/10.1111/ele.12382>

740 Kitamura, K., Nakai, Y., Suzuki, S., Ohtani, Y., Yamanoi, K., Sakamoto, T., 2012.

741 Interannual variability of net ecosystem production for a broadleaf deciduous

742 forest in Sapporo, northern Japan. *J. For. Res.* 17, 323–332.

743 <https://doi.org/10.1007/s10310-012-0335-4>

744 Knohl, A., Schulze, E., Kolle, O., Buchmann, N., 2003. Large carbon uptake by an

745 unmanaged 250-year-old deciduous forest in Central Germany 118, 151–167.

746 [https://doi.org/10.1016/S0168-1923\(03\)00115-1](https://doi.org/10.1016/S0168-1923(03)00115-1)

747 Kolle, O., Rebmann, C., 2010. 10: EddySoft : documentation of a software package to

748 acquire and process eddy covariance data. Tech. reports, Max-Planck-Institut für

749 Biogeochem. 10.

750 Law, B., Falge, E., Gu, L., Baldocchi, D., Bakwin, P., Berbigier, P., Davis, K., Dolman, A.,

751 Falk, M., Fuentes, J., Goldstein, A., Granier, A., Grelle, A., Hollinger, D., Janssens, I.,

752 Jarvis, P., Jensen, N., Katul, G., Mahli, Y., Matteucci, G., Meyers, T., Monson, R.,

753 Munger, W., Oechel, W., Olson, R., Pilegaard, K., Paw U, K., Thorgeirsson, H.,

754 Valentini, R., Verma, S., Vesala, T., Wilson, K., Wofsy, S., 2002. Environmental

755 controls over carbon dioxide and water vapor exchange of terrestrial vegetation.

756 *Agric. For. Meteorol.* 113, 97–120. [https://doi.org/10.1016/S0168-1923\(02\)00104-1](https://doi.org/10.1016/S0168-1923(02)00104-1)

757 Lenth, R., Love, J., 2017. Package “lsmeans” Title Least-Squares Means.

758 <https://doi.org/10.1080/00031305.1980.10483031>

759 Luo, Y., Keenan, T.F., Smith, M., 2015. Predictability of the terrestrial carbon cycle. *Glob.*

760 *Chang. Biol.* 21, 1737–1751. <https://doi.org/10.1111/gcb.12766>

761 Luyssaert, S., 2014. Land management and land-cover change have impacts of similar

762 magnitude on surface temperature. *Nat. Clim. Chang.* 4, 5.

763 <https://doi.org/10.1038/NCLIMATE2196>

764 Ma, S., Baldocchi, D.D., Xu, L., Hehn, T., 2007. Inter-annual variability in carbon dioxide

765 exchange of an oak/grass savanna and open grassland in California. *Agric. For.*

766 *Meteorol.* 147, 157–171. <https://doi.org/10.1016/j.agrformet.2007.07.008>

767 Mund, M., 2004. Carbon pools of European beech forests (*Fagus sylvatica*) under

768 different silvicultural management 189, 256.

769 Mund, M., Kutsch, W.L., Wirth, C., Kahl, T., Knohl, A., Skomarkova, M. V., Schulze, E.D.,

770 2010. The influence of climate and fructification on the inter-annual variability of

771 stem growth and net primary productivity in an old-growth, mixed beech forest.

772 *Tree Physiol.* 30, 689–704. <https://doi.org/10.1093/treephys/tpq027>

773 Musavi, T., Migliavacca, M., Reichstein, M., Kattge, J., Wirth, C., Black, T.A., Janssens, I.,

774 Knohl, A., Loustau, D., Rouspard, O., Varlagin, A., Rambal, S., Cescatti, A., Gianelle,

775 D., Kondo, H., Tamrakar, R., Mahecha, M.D., 2017. Stand age and species richness

776 dampen interannual variation of ecosystem-level photosynthetic capacity. *Nat.*

777 *Ecol. Evol.* 1, 1–6. <https://doi.org/10.1038/s41559-016-0048>

778 Nathans, L.L., Oswald, F.L., Nimon, K., 2012. Interpreting Multiple Linear Regression: A

779 Guidebook of Variable Importance. *Pract. Assessment, Res. Eval.* 17, 1–19.

780 <https://doi.org/10.3102/00346543074004525>

781 Novick, K.A., Oishi, A.C., Ward, E.J., Siqueira, M.B.S., Juang, J.-Y., Stoy, P.C., 2015. On the

782 difference in the net ecosystem exchange of CO₂ between deciduous and

783 evergreen forests in the southeastern United States. *Glob. Chang. Biol.* 21, 827–

784 842. <https://doi.org/10.1111/gcb.12723>

785 Obeso, J.R., 2002. The costs of reproduction in plants *Author. New Phytol.* 155, 321–348.

786 <https://doi.org/10.1046/j.1469-8137.2002.00477.x>

787 Pereira, J.S., Mateus, J.A., Aires, L.M., Pita, G., Pio, C., David, J.S., Andrade, V., Banza, J.,
 788 David, T.S., Paço, T.A., Rodrigues, A., 2007. Net ecosystem carbon exchange in
 789 three contrasting Mediterranean ecosystems – the effect of drought.
 790 Biogeosciences 4, 791–802. <https://doi.org/10.5194/bg-4-791-2007>
 791 Pilegaard, K., Ibrom, A., Courtney, M.S., Hummelshøj, P., Jensen, N.O., 2011. Increasing
 792 net CO₂ uptake by a Danish beech forest during the period from 1996 to 2009.
 793 Agric. For. Meteorol. 151, 934–946.
 794 <https://doi.org/10.1016/J.AGRFORMET.2011.02.013>
 795 R Core Team, 2017. R: A Language and Environment for Statistical Computing.
 796 Reichstein, M., Falge, E., Baldocchi, D., Papale, D., Aubinet, M., Berbigier, P., Bernhofer,
 797 C., Buchmann, N., Gilmanov, T., Granier, A., Grunwald, T., Havrahkova, K.,
 798 Ilvesniemi, H., Janous, D., Knohl, A., Laurila, T., Lohila, A., Loustau, D., Matteucci, G.,
 799 Meyers, T., Miglietta, F., Ourcival, J., Pumpanen, J., Rambal, S., Rotenberg, E., Sanz,
 800 M., Tenhunen, J., Seufert, G., Vaccari, F., Vesala, T., Yakir, C., Valentini, R., 2005. On
 801 the separation of net ecosystem exchange into assimilation and ecosystem
 802 respiration: review and improved algorithm. Glob. Chang. Biol. 11, 1424–1439.
 803 Richardson, A.D., Hollinger, D.Y., Dail, D.B., Lee, J.T., Munger, J.W., O’Keefe, J., 2009.
 804 Influence of spring phenology on seasonal and annual carbon balance in two
 805 contrasting New England forests. Tree Physiol. 29, 321–331.
 806 <https://doi.org/10.1093/treephys/tpn040>
 807 Sala, A., Hopping, K., McIntire, E.J.B., Delzon, S., Crone, E.E., 2012. Masting in whitebark
 808 pine (*Pinus albicaulis*) depletes stored nutrients. New Phytol. 196, 189–199.
 809 <https://doi.org/10.1111/j.1469-8137.2012.04257.x>
 810 Selås, V., Piovesan, G., Adams, J.M., Bernabei, M., 2002. Climatic factors controlling
 811 reproduction and growth of Norway spruce in southern Norway. Can. J. For. Res.
 812 32, 217–225. <https://doi.org/10.1139/x01-192>
 813 Shao, J., Zhou, X., He, H., Yu, G., Wang, H., Luo, Y., Chen, J., Gu, L., Li, B., 2014.
 814 Partitioning Climatic and Biotic Effects on Interannual Variability of Ecosystem
 815 Carbon Exchange in Three Ecosystems. Ecosystems 17, 1186–1201.
 816 <https://doi.org/10.1007/s10021-014-9786-0>
 817 Shao, J., Zhou, X., Luo, Y., Li, B., Aurela, M., Billesbach, D., Blanken, P.D., Bracho, R.,
 818 Chen, J., Fischer, M., Fu, Y., Gu, L., Han, S., He, Y., Kolb, T., Li, Y., Nagy, Z., Niu, S.,
 819 Oechel, W.C., Pinter, K., Shi, P., Suyker, A., Torn, M., Varlagin, A., Wang, H., Yan, J.,
 820 Yu, G., Zhang, J., 2016. Direct and indirect effects of climatic variations on the
 821 interannual variability in net ecosystem exchange across terrestrial ecosystems.
 822 Tellus B Chem. Phys. Meteorol. 68, 30575.
 823 <https://doi.org/10.3402/tellusb.v68.30575>
 824 Shao, J., Zhou, X., Luo, Y., Li, B., Aurela, M., Billesbach, D., Blanken, P.D., Bracho, R.,
 825 Chen, J., Fischer, M., Fu, Y., Gu, L., Han, S., He, Y., Kolb, T., Li, Y., Nagy, Z., Niu, S.,
 826 Oechel, W.C., Pinter, K., Shi, P., Suyker, A., Torn, M., Varlagin, A., Wang, H., Yan, J.,
 827 Yu, G., Zhang, J., 2015. Biotic and climatic controls on interannual variability in
 828 carbon fluxes across terrestrial ecosystems. Agric. For. Meteorol. 205, 11–22.
 829 <https://doi.org/10.1016/j.agrformet.2015.02.007>
 830 Urbanski, S., Barford, C., Wofsy, S., Kucharik, C., Pyle, E., Budney, J., McKain, K.,
 831 Fitzjarrald, D., Czikowsky, M., Munger, J.W., 2007. Factors controlling CO₂
 832 exchange on timescales from hourly to decadal at Harvard Forest. J. Geophys. Res.
 833 Biogeosciences 112, 1–25. <https://doi.org/10.1029/2006JG000293>
 834 Wilkinson, M., Eaton, E.L., Broadmeadow, M.S.J., Morison, J.I.L., 2012. Inter-annual

variation of carbon uptake by a plantation oak woodland in south-eastern England. *Biogeosciences* 9, 5373–5389. <https://doi.org/10.5194/bg-9-5373-2012>


Wilson, K.B., Baldocchi, D.D., 2001. Comparing independent estimates of carbon dioxide exchange over 5 years at a deciduous forest in the southeastern United States. *J. Geophys. Res. Atmos.* 106, 34167–34178. <https://doi.org/10.1029/2001jd000624>

Wu, C., Chen, J.M., Black, T.A., Price, D.T., Kurz, W.A., Desai, A.R., Gonsamo, A., Jassal, R.S., Gough, C.M., Bohrer, G., Dragoni, D., Herbst, M., Gielen, B., Berninger, F., Vesala, T., Mammarella, I., Pilegaard, K., Blanken, P.D., 2013. Interannual variability of net ecosystem productivity in forests is explained by carbon flux phenology in autumn. *Glob. Ecol. Biogeogr.* 22, 994–1006. <https://doi.org/10.1111/geb.12044>


Wu, C., Chen, J.M., Gonsamo, A., Price, D.T., Black, T.A., Kurz, W.A., 2012. Interannual variability of net carbon exchange is related to the lag between the end-dates of net carbon uptake and photosynthesis: Evidence from long records at two contrasting forest stands. *Agric. For. Meteorol.* 164, 29–38. <https://doi.org/10.1016/j.agrformet.2012.05.002>

Wutzler, T., Reichstein, M., Moffat, A.M., Menzer, O., Migliavacca, M., Sickel, K., Sigut, L., 2018. Title Post Processing of (Half-)Hourly Eddy-Covariance Measurements.

Yuan, W., Luo, Y., Richardson, A.D., Oren, R., Luyssaert, S., Janssens, I.A., Ceulemans, R., Zhou, X., Grünwald, T., Aubinet, M., Berhofer, C., Baldocchi, D.D., Chen, J., Dunn, A.L., Deforest, J.L., Dragoni, D., Goldstein, A.H., Moors, E., Munger, J.W., Monson, R.K., Suyker, A.E., Starr, G., Scott, R.L., Tenhunen, J., Verma, S.B., Vesala, T., Wofsy, S.T.E., 2009. Latitudinal patterns of magnitude and interannual variability in net ecosystem exchange regulated by biological and environmental variables. *Glob. Chang. Biol.* 15, 2905–2920. <https://doi.org/10.1111/j.1365-2486.2009.01870.x>



862
863 Figure S1: Annual footprint weighted thinning in eddy flux footprint area of DE-Lnf from
864 1995 to 2016. The highest thinning of 998 g C m⁻² was conducted in 2005 from the
865 footprint area.



866
867 Figure S2: Linear regression between mean annual T_{soil} of DE-Lnf and DE-Hai. Dashed
868 line represents a line with slope 1. Systematic difference between sites is due to
869 difference in measurement depths (5 vs 4 cm in DE-Hai and DE-Lnf, respectively).
870

Table S1: Average Annual values (mean and sd) and the temporal trend during the study period for all dependent and selected predictor variables for both sites. Units for GSS (growing season start day), and GSE (growing season end day) is doy yr⁻¹ where doy indicates Julian day of the year. Trend is the temporal trend and * indicates the statistically significant values at 5% significance level.

| Variables | Units | DE-Hai | | | DE-Lnf | | |
|-------------------|--------------------------------------|--------|------|-------|--------|-------|--------------|
| | | mean | sd | trend | mean | sd | trend |
| R _g | W m ⁻² | 122 | 6.5 | 0.18 | 124.0 | 7.2 | 0.62 |
| T _{air} | °C yr ⁻¹ | 8.34 | 0.72 | 0.03 | 8.30 | 0.70 | 0.05 |
| T _{soil} | °C yr ⁻¹ | 7.61 | 0.36 | 0.01 | 8.23 | 0.38 | 0.02 |
| VPD | hPa yr ⁻¹ | 3.45 | 0.56 | 0.02 | 3.28 | 0.48 | 0 |
| Rain | mm yr ⁻¹ | 744 | 152 | -6.76 | 601 | 154 | -9.12 |
| WAI | yr ⁻¹ | 0.86 | 0.09 | -0.01 | 0.84 | 0.1 | 0 |
| NEP | g C m ⁻² yr ⁻¹ | 487 | 57.8 | -3.14 | 585 | 144 | 21.8* |
| GPP | g C m ⁻² yr ⁻¹ | 1558 | 118 | 4.25 | 1627 | 164 | 25.5* |
| Reco | g C m ⁻² yr ⁻¹ | 1071 | 96 | 0.28 | 1042 | 60 | 0.08 |
| GSS | doy yr ⁻¹ | 125 | 7 | -0.42 | 121 | 5 | -0.29 |
| GSE | doy yr ⁻¹ | 289 | 4 | -0.35 | 288 | 5 | 0.45 |
| GSL | day yr ⁻¹ | 164 | 7 | 0.07 | 168 | 6 | 0.75* |
| Fruit production | g C m ⁻² yr ⁻¹ | 73.2 | 77.9 | 4.53 | 91.2 | 113.1 | 5.43 |

Table S2: Growing season start day (GSS), end day (GSE) and length (GSL) for DE-Hai and DE-Lnf during the study period.

| Year | DE-Hai | | | DE-Lnf | | |
|---------|-----------|-----------|------------|-----------|-----------|------------|
| | GSS (doy) | GSE (doy) | GSL (days) | GSS (doy) | GSE (doy) | GSL (days) |
| 2003 | 125 | 292 | 167 | 123 | 290 | 167 |
| 2004 | 123 | 288 | 165 | 121 | 281 | 160 |
| 2005 | 125 | 293 | 168 | 122 | 291 | 169 |
| 2006 | 127 | 294 | 167 | 126 | 294 | 168 |
| 2010 | 134 | 283 | 149 | 123 | 283 | 160 |
| 2011 | 117 | 288 | 171 | 114 | 285 | 171 |
| 2012 | 129 | 289 | 160 | 124 | 290 | 166 |
| 2013 | 131 | 288 | 157 | 130 | 292 | 162 |
| 2014 | 111 | 284 | 173 | 112 | 288 | 176 |
| 2015 | 114 | 284 | 170 | 117 | 293 | 176 |
| 2016 | 130 | 292 | 162 | 117 | 292 | 175 |
| Average | 125 | 289 | 164 | 121 | 289 | 168 |
| Sd | 7 | 4 | 7 | 5 | 4 | 6 |




Figure S3: Relationship between T_{air} and T_{soil} with growing season matrix. GSS is growing season start day indicated by day of the year (doy), GSE is growing season end day indicated by day of the year (doy) and GSL is growing season length indicated by number of days. The solid lines represent linear regression lines and dashed lines its confidence interval at 5% significance level. *** indicate statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.




Figure S4: Simple linear regression between NEP, GPP and Reco with growing season start day (GSS), growing season end day (GSE) and growing season length day (GSL). The solid lines represent linear regression lines and dashed lines confidence interval at 5% significance level. *** indicate statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.

Table S3: Mean annual fluxes for two periods (2003-2006 and 2010-2016) and differences between two sites. Differences between sites were tested using paired t-test.

| period | Flux | DE-Hai (mean \pm sd) [g C m ⁻² yr ⁻¹] | DE-Lnf (mean \pm sd) [g C m ⁻² yr ⁻¹] | Diff _(DE-Lnf-DE-Hai) [g C m ⁻² yr ⁻¹] |
|-----------|------|---|---|--|
| 2003-2006 | NEP | 506 \pm 84 | 476 \pm 122 | 30 |
| | GPP | 1542 \pm 76 | 1517 \pm 128 | 25 |
| | Reco | 1035 \pm 50 | 1040 \pm 25 | 2 |
| 2010-2016 | NEP | 476 \pm 41 | 647 \pm 121 | 171** |
| | GPP | 1568 \pm 142 | 1689 \pm 156 | 121** |
| | Reco | 1092 \pm 112 | 1042 \pm 76 | -50 |

Table S4: Difference between the two sites with respect to slopes between each of the three significant driving variables (T_{soil} , FP, time) and the three flux quantities (NEP,

GPP, Reco) in terms of the residual variance remaining after the effects of the other two driving variables (e.g. FP and time, in the case of T_{soil}) have been removed.

| CO ₂ flux | Variable | Slope difference (DE-Hai – DE-Lnf) | SE |
|----------------------|------------|---------------------------------------|------|
| NEP | T_{soil} | 32.6 | 44.5 |
| GPP | T_{soil} | 26.6 | 79.6 |
| Reco | T_{soil} | -5.9 | 63.4 |
| NEP | FP | 0.4* | 0.2 |
| GPP | FP | 0.3 | 0.3 |
| Reco | FP | -0.1 | 0.3 |
| NEP | Time | -21.2** | 4.3 |
| GPP | Time | -11.9(*) | 7.0 |
| Reco | Time | 9.3(*) | 5.5 |

Slope difference indicates the difference between two sites with respect to slopes between each of the three significant driving variables (T_{soil} , FP, time) and the three flux quantities (NEP, GPP, Reco) in terms of the residual variance remaining after the effects of the other two driving variables (e.g. FP and time, in the case of T_{soil}) have been removed. SE is the standard error of that difference. *** indicate statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.




Figure S5: Relationship of growing season start day (GSS) and growing season length (GSL) with spring air (AM T_{air}) and soil temperature (AM T_{soil}). The solid lines represent linear regressions and dotted lines 95% confidence interval. *** indicates statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.




Figure S6: Regression between annual NEP and annual mean of meteorological variables (except for rain which was summed). The solid lines represent linear regressions and dotted lines 95% confidence interval. *** indicates statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.




Figure S7: Regression between annual GPP and annual mean of meteorological variables (except for rain which was summed). The solid lines represent linear regressions and dotted lines 95% confidence interval. *** indicates statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.




Figure S8: Linear regression between residual of NEP, GPP and Reco after removing the effect of time and fruit production (FP). CI is 95% confidence interval of slope and rsq the coefficient of determination of linear regression, and sig its significance. The solid lines represent linear regression lines and dashed lines confidence interval at 5% significance level. *** indicate statistical significance at $p < 0.001$; ** significant at $p < 0.01$; * significant at $p < 0.05$; and (*) significant at $p < 0.1$.

928 Table S5: Temperate broad-leaved deciduous forests with long term eddy covariance measurements. The table was modified after Baldocchi et al.
 929 (2018). CV is coefficient of variation (standard deviation divided by mean).

| Site | Country | n | NEP | | | GPP | | | Reco | | | Reference |
|---------------------------|-------------------|----|---|---|----------|---|---|----------|---|---|----------|--|
| | | | mean (g C m ⁻² yr ⁻¹) | sd (g C m ⁻² yr ⁻¹) | CV -- | mean (g C m ⁻² yr ⁻¹) | sd (g C m ⁻² yr ⁻¹) | CV -- | mean (g C m ⁻² yr ⁻¹) | sd (g C m ⁻² yr ⁻¹) | CV -- | |
| Borden | Canada-Ontario | 18 | 177 | 116 | 0.66 | 1373 | 164 | 0.12 | 1196 | 188 | 0.16 | Froelich et al. (2015) Pilegaard et al. (2011) Granier et al. (2008) |
| Soroe | Denmark | 13 | 156 | 103 | 0.66 | 1727 | 136 | 0.08 | 1570 | 97 | 0.06 | |
| Hesse | France | 10 | 386 | 171 | 0.44 | 1397 | 192 | 0.14 | 1011 | 137 | 0.14 | |
| Takayama | Japan | 9 | 237 | 98 | 0.41 | 1110 | 409 | 0.37 | 829 | 264 | 0.32 | Wilkinson et al. (2012) Sulman et al. (2016) Urbanski et al. (2007) |
| Straights Inclosure | United Kingdom | 12 | 486 | 115 | 0.24 | 1993 | 275 | 0.14 | 1548 | 192 | 0.12 | |
| Morgan-Monroe | US-Indiana | 13 | 351 | 81 | 0.23 | 1452 | 118 | 0.08 | 1098 | 82 | 0.07 | |
| Harvard Forest, Petersham | US-Massachusetts | 13 | 245 | 100 | 0.41 | 1400 | 164 | 0.12 | 1153 | 105 | 0.09 | Shao et al. (2014) Novick et al. (2015) Wilson and Baldocchi (2001) |
| Ozarks | US-Missouri | 5 | 479 | 65 | 0.14 | 1125 | 164 | 0.15 | 646 | 121 | 0.19 | |
| Duke Forest, Durham | US-North Carolina | 8 | 402 | 96 | 0.24 | 1982 | 300 | 0.15 | 1580 | 237 | 0.15 | |
| Oak Ridge | US-Tennessee | 5 | 577 | 63 | 0.11 | Na | Na | Na | Na | Na | Na | |
| Average | | | 350 | 100 | 0.35 | 1506 | 214 | 0.15 | 1181 | 158 | 0.14 | |